

Anbesol Gel Maximum Strength

<p>Drug Facts</p> <table style="width: 100%; border: none;"> <tr> <td style="width: 50%;">Active ingredient</td> <td style="width: 50%;">Purpose</td> </tr> <tr> <td>Benzocaine 20%</td> <td>Oral pain reliever</td> </tr> </table> <p>Uses</p> <ul style="list-style-type: none"> ■ temporarily relieves pain associated with the following mouth and gum irritations: <ul style="list-style-type: none"> ■ toothache ■ canker sores ■ minor dental procedures ■ dentures ■ sore gums ■ braces <p>Warnings</p> <p>METHEMOGLOBINEMIA WARNING: Use of this product may cause methemoglobinemia, a rare but serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:</p> <ul style="list-style-type: none"> ■ pale, gray, or blue colored skin (cyanosis) ■ headache ■ rapid heart rate ■ shortness of breath ■ dizziness or lightheadedness ■ fatigue or lack of energy <p>Allergy alert: Do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics.</p> <p>Do not use</p> <ul style="list-style-type: none"> ■ for teething ■ in children under 2 years of age <p>When using this product</p> <ul style="list-style-type: none"> ■ avoid contact with the eyes ■ do not exceed recommended dosage ■ do not use for more than 7 days unless directed by a doctor/dentist <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ■ sore mouth symptoms do not improve in 7 days ■ irritation, pain, or redness lasts or worsens ■ swelling, rash, or fever develops 	Active ingredient	Purpose	Benzocaine 20%	Oral pain reliever	<p>Drug Facts (continued)</p> <p>Keep out of reach of children. If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away.</p> <p>Directions</p> <ul style="list-style-type: none"> ■ to open tube, cut tip of the tube on score mark with scissors <table style="width: 100%; border: none;"> <tr> <td style="width: 50%;">adults and children 2 years of age and over</td> <td style="width: 50%;">apply a pea-size amount to the affected area up to 4 times daily or as directed by a doctor/dentist</td> </tr> <tr> <td>children under 12 years of age</td> <td>adult supervision should be given in the use of this product</td> </tr> <tr> <td>children under 2 years of age</td> <td>do not use</td> </tr> </table> <ul style="list-style-type: none"> ■ for denture irritation: <ul style="list-style-type: none"> ■ apply thin layer to the affected area ■ do not reinsert dental work until irritation/pain is relieved ■ rinse mouth well before reinserting <p>Other information</p> <ul style="list-style-type: none"> ■ store at 20-25°C (68-77°F) ■ do not refrigerate <p>Inactive ingredients benzyl alcohol, carbomer 934P, D&C yellow no. 10, FD&C blue no. 1, FD&C red no. 40, glycerin, methylparaben, natural flavor, polyethylene glycol, polysorbate 80, propylene glycol, saccharin</p> <p>Questions or comments? Call 1-888-594-0673 weekdays 9 AM to 5 PM EST</p>	adults and children 2 years of age and over	apply a pea-size amount to the affected area up to 4 times daily or as directed by a doctor/dentist	children under 12 years of age	adult supervision should be given in the use of this product	children under 2 years of age	do not use
Active ingredient	Purpose										
Benzocaine 20%	Oral pain reliever										
adults and children 2 years of age and over	apply a pea-size amount to the affected area up to 4 times daily or as directed by a doctor/dentist										
children under 12 years of age	adult supervision should be given in the use of this product										
children under 2 years of age	do not use										