

Anbesol Liquid Maximum Strength

Drug Facts		Drug Facts (continued)	
Active ingredient Benzocaine 20%		Purpose Oral pain reliever	
Uses ■ temporarily relieves pain associated with the following mouth and gum irritations: ■ toothache ■ sore gums ■ canker sores ■ braces ■ minor dental procedures ■ dentures		Directions adults and children 2 years of age and over wipe liquid on with cotton, or cotton swab, or fingertip, apply to the affected area up to 4 times daily or as directed by a doctor/dentist children under 12 years of age adult supervision should be given in the use of this product children under 2 years of age do not use ■ for denture irritation: ■ apply thin layer to the affected area ■ do not reinsert dental work until irritation/pain is relieved ■ rinse mouth well before reinserting	
Warnings METHEMOGLOBINEMIA WARNING: Use of this product may cause methemoglobinemia, a rare but serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops: ■ pale, gray, or blue colored skin (cyanosis) ■ headache ■ rapid heart rate ■ shortness of breath ■ dizziness or lightheadedness ■ fatigue or lack of energy Allergy alert: Do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics.		Other information store at 20-25°C (68-77°F)	
Do not use ■ for teething ■ in children under 2 years of age		Inactive ingredients benzyl alcohol, D&C yellow no. 10, FD&C blue no. 1, FD&C red no. 40, methylparaben, natural flavor, polyethylene glycol, polysorbate 80, propylene glycol, saccharin	
When using this product ■ avoid contact with the eyes ■ do not exceed recommended dosage ■ do not use for more than 7 days unless directed by a doctor/dentist		Questions or comments? Call 1-888-594-0673 weekdays 9 AM to 5 PM EST	
Stop use and ask a doctor if ■ sore mouth symptoms do not improve in 7 days ■ irritation, pain, or redness lasts or worsens ■ swelling, rash, or fever develops			